


# Tabulka mísitelnosti biopreparátů

	Fungicidy a pomocné prostředky								
	Myco-Sin	Altela, Memcomba	HF - Mycol	Aqua Vitrin K	VitiSan	Kumulus WG	Síra BL	Flowbrix	Cocana
Myco-Sin			!	!	!	!	!	!	!
Altela, Memcomba									
HF - Mycol	!								
Aqua Vitrin K	!								
VitiSan	!								
Kumulus WG	!								
Síra BL	!								
Flowbrix	!								
Cocana	!								
WETCIT							*	*	
PREV-B2							*	*	
VermiFit A, VermiFit B				!					
ProKton				!			!	!	
Hycol-E víno				!			!		
polysulfid vápenatý	!		!	!	!		!	!	!
Lepinox Plus				!	!	!	!	!	!
Madex, Madex TOP	!			!		?	!	!	
NeemAzal -T/S	!		?	?	?	?	**	!	
Isonet, Isomate									
<i>Typhlodromus pyri</i>							!		!

 kombinaci preparátů DOPORUČUJEME


 při vyšší koncentraci je MÍRNĚ TOXICKÝ vůči dravému roztoči

 kombinace zatím nebyla testována

 není důvod uvažovat o míchání

 Toxicita ostatních přípravků používaných v révě vinné a v jabloních je uvedena na webových stránkách [www.biocont.cz](http://www.biocont.cz)

Smáždědla	Hnojiva						Insekticidy				Hodnota pH	
	WETCIT	PREV-B2	VermiFit A, VermiFit B	ProKton	Hycol-E víno	polysulfid vápenatý	Lepinox Plus	Madex, Madex TOP	NeemAzal -T/S	Isonet, Isomate		<i>Typhlodromus pyri</i>
						!		!	!			3,8
												5–6
						!			?			8,2
			!	!	!	!	!	!	?			10,5
						!	!		?			8,1–8,4
							!		?			6–7
	*	*			!						!	6–7
	*	*	!	!	!	!	!	!	**			8,6
			!		!	!	!	!	!		!	12,5
					!							8,2–8,5
					!							8,2–8,5
					!							6–7
					!							6–7
												5,5–7
	!	!	!	!			!	!	!		!	11
						!						4–6,5
						!						6–7
						!						7
												-
						!						-

 koncentrace je buď nevhodná (přípravky vzájemně reagují) nebo fytotoxická – NELZE MÍCHAT

\* v případě velmi vysokých teplot nad 25 °C použijte nižší koncentraci a s produktem PREV-B2 nemísíte.

\*\* lze mísit s nízkými dávkami Kocide 2000 (tzn. do 500 g/ha, tj. cca 0,1 %)