

- [1] Schwarze H, Küntscher M, Uhlig C, Hierlemann H, Prantl L, Ottomann C, et al. Suprathel, a New Skin Substitute, in the Management of Partial- Thickness Wounds. *Ann Plast Surg* 2008;60:181–5. doi:10.1097/SAP.0b013e318056bbf6.
- [2] Uhlig C, Rapp M, Dittel KK. Neue strategien zur behandlung thermisch geschädigter hände unter berücksichtigung des epithelersatzes Suprathel®. *Handchirurgie Mikrochirurgie Plast Chir* 2007;39:314–9. doi:10.1055/s-2007-965234.
- [3] Uhlig C, Rapp M, Hartmann B, Hierlemann H, Planck H, Dittel K-KK, et al. Suprathel-an innovative, resorbable skin substitute for the treatment of burn victims. *Burns* 2007;33:221–9. doi:10.1016/j.burns.2006.04.024.
- [4] Hardy Schwarze, MD,† Markus Küntscher, MD,† Christian Uhlig, MD,\* Helmut Hierlemann, MD,‡ Lukas Prantl, MD,† Christian Ottomann, MD,† and Bernd Hartmann M, Schwarze H, Küntscher M, Uhlig C, Hierlemann H, Prantl L, et al. Suprathel, a new skin substitute, in the management of partial-thickness burn wounds: results of a clinical study. *Ann Plast Surg* 2008;60:181–5. doi:10.1097/SAP.0b013e318056bbf6.
- [5] Blome-Eberwein SAM, Patrick Pagella RN C, Deborah Boorse RN C, MD HA. Results from Application of an Absorbable Synthetic Membrane to

- Superficial and Deep Second Degree Burn Wounds. Boston: 2014.
- [6] Glik J, Kawecki M, Kitala D, Kłama-Baryła A, Łabuś W, Grabowski M, et al. A new option for definitive burn wound closure – pair matching type of retrospective case–control study of hand burns in the hospitalised patients group in the Dr Stanisław Sakiel Centre for Burn Treatment between 2009 and 2015. *Int Wound J* 2017;14:849–55. doi:10.1111/iwj.12720.
- [7] Held M, Rothenberger J, Engelke AS, Tolzmann DS, Esfahani BJ, Schaller HE, et al. Evaluation of commonly used temporary skin dressings and a newly developed collagen matrix for treatment of superficial wounds. *Adv Ski Wound Care* 2015;28:551–4. doi:10.1097/01.ASW.0000473136.66014.69.
- [8] Kaartinen IS, V?lisuo PO, Alander JT, Kuokkanen HO. Objective scar assessment - A new method using standardized digital imaging and spectral modelling. *Burns* 2011;37:74–81. doi:10.1016/j.burns.2010.03.008.
- [9] Behr B, Megerle KO, Germann G, Kloeters O, Cermann G, Kloeters O. New concepts in local burn wound therapy. *Handchirurgie Mikrochirurgie Plast Chir* 2008;40:361–6. doi:10.1055/s-2008-1039062.
- [10] Lampe KJ, Namba RM, Silverman TR, Bjugstad KB. Impact of Lactic Acid on Cell Proliferation and Free Radical Induced Cell Death in Monolayer

- Cultures of Neural Precursor Cells. *Biotechnol Bioeng* 2009;103:1214–23. doi:10.1002/bit.22352. Impact.
- [11] Herz H, Blake DR, Grootveld M. Multicomponent investigations of the hydrogen peroxide- and hydroxyl radical-scavenging antioxidant capacities of biofluids: The roles of endogenous pyruvate and lactate. *Free Radic Res* 1997;26:19–35. doi:10.3109/10715769709097781.
- [12] Philp A, Macdonald AL, Watt PW. Lactate—a signal coordinating cell and systemic function. *J Exp Biol* 2005;208:4561–75. doi:10.1242/jeb.01961.
- [13] Miller BF, Fattor JA, Jacobs KA, Horning MA, Navazio F, Lindinger MI, et al. Lactate and glucose interactions during rest and exercise in men: Effect of exogenous lactate infusion. *J Physiol* 2002;544:963–75. doi:10.1113/jphysiol.2002.027128.
- [14] YAMADA H, IWAKI Y, KITAOKA R, FUJITANI M, SHIBAKUSA T, FUJIKAWA T, et al. Blood Lactate Functions as a Signal for Enhancing Fatty Acid Metabolism during Exercise via TGF- $\beta$  in the Brain. *J Nutr Sci Vitaminol (Tokyo)* 2012;58:88–95. doi:10.3177/jnsv.58.88.
- [15] Chappell D, Hofmann-Kiefer K, Jacob M, Rehm M, Briegel J, Welsch U, et al. TNF- $\alpha$  induced shedding of the endothelial glykokalyx is prevented by hydrocortisone and antithrombin. *Basic Res Cardiol* 2009;104:78–89. doi:10.1007/s00395-008-0749-5.

- [16] Kottmann RM, Trawick E, Judge JL, Wahl LA, Epa AP, Owens KM, et al. Pharmacologic inhibition of lactate production prevents myofibroblast differentiation. *Am J Physiol Cell Mol Physiol* 2015;309:L1305–12. doi:10.1152/ajplung.00058.2015.
- [17] SILVIA WAGNER, PhD; M. ZAMIRUL HUSSAIN, PhD; THOMAS K. HUNT MdBacHD, BECKER Md, Wagner S, Hussain MZ, Hunt TK, Bacic BBBBB, et al. Stimulation of fibroblast proliferation by lactate-mediated oxidants. *Wound Repair Regen* 2004;12:368–73. doi:10.1111/j.1067-1927.2004.012315.x.
- [18] Trabold O, Wagner S, Wicke C, Scheuenstuhl H, Hussain MZ, Rosen N, et al. Lactate and oxygen constitute a fundamental regulatory mechanism in wound healing. *Wound Repair Regen* 2003;11:504–9. doi:10.1046/j.1524-475X.2003.11621.x.
- [19] McMahon S, Charbonneau M, Grandmont S, Richard DE, Dubois CM. Transforming growth factor  $\beta$ 1 induces hypoxia-inducible factor-1 stabilization through selective inhibition of PHD2 expression. *J Biol Chem* 2006;281:24171–81. doi:10.1074/jbc.M604507200.
- [20] Constant JS1, Feng JJ, Zabel DD, Yuan H, Suh DY, Scheuenstuhl H, Hunt TK HM, Constant JS, Feng JJ, Zabel DD, Yuan H, Suh DY, et al. Lactate elicits vascular endothelial growth factor from macrophages: a possible alternative to hypoxia. *Wound Repair Regen*

- 2000;8:353–60.  
doi:10.1046/j.1524-475X.2000.00353.x.
- [21] Rousselle P, Braye F, Dayan G. Re-epithelialization of adult skin wounds: Cellular mechanisms and therapeutic strategies. *Adv Drug Deliv Rev* 2018. doi:10.1016/j.addr.2018.06.019.
- [22] Sun T, Adra S, Smallwood R, Holcombe M, MacNeil S. Exploring hypotheses of the actions of TGF- $\beta$ 1 in epidermal wound healing using a 3D computational multiscale model of the human epidermis. *PLoS One* 2009;4. doi:10.1371/journal.pone.0008515.
- [23] Wang P, Liu X, Xu P, Lu J, Wang R, Mu W. Decorin reduces hypertrophic scarring through inhibition of the TGF- $\beta$ 1/Smad signaling pathway in a rat osteomyelitis model. *Exp Ther Med* 2016;12:2102–8. doi:10.3892/etm.2016.3591.
- [24] Phan TT, See P, Lee ST, Chan SY. Protective effects of curcumin against oxidative damage on skin cel

Is in vitro: its implication for wound healing. J Trauma 2001;51:927–31. doi:10.1097/00005373-200111000-00017.

[25] Wahl P, Bloch W, Mester J. Moderne Betrachtungsweisen des Laktats: Laktat ein ueberschaetztes und zugleich unterschaeztetes Molekuehl. Schweizerische Zeitschrift Fur Sport Und Sport 2009;57:100–7.

[26] Brooks GA, Brooks TG, Brooks S. Laktat als metabolisches signal der genexpression. Dtsch Z Sportmed 2008;59:280–6.

[27] Gürünlüoğlu K, Demircan M, Taşçi A, Üremiş MM, Türköz Y, Bağ HG, et al. The effects of two different burn dressings on serum oxidativ

e stress indicators in children with partial burn. J

Burn Care Res 2017;00:1–19. doi:10.1080/13632469.2017.1342302.

[28] Gürünlüoğlu K, Demircan M, Koç A, Koçbiyik A, Taşçi A, Durmuş K, et al. The effects of different burn dressings on length of telomere and expression of telomerase in children with thermal burns. J Burn Care Res 2019. doi:10.1093/jbcr/irz019.

[29] E. Schriek; M. Sinnig. The Use of Caprolactone Dressing in Pediatric Burns - A Gold Standard ? ABA 2018, 2018.

[30] Blome-Eberwein SA, Amani H, Lozano D, Gogal C. Second-Degree Burn Care with a Lactic Acid Based Biodegradable Skin Substitute in 229 Pediatric and Adult Patients. J Burn Care Res 2018;39:S223–S223.

doi:10.1093/jbcr/iry0  
06.423.

- [31] Rothenberger J, Held  
M, Stolz A, Tschumi  
C, Olariu R, Pdf D.  
Use of a Polylactide -  
based Copolymer as a  
Temporary Skin  
Substitute for a  
Patient With Moist  
Desquamation Due to  
Radiation. Wounds  
2016;7:1E26-30.