

## **Patients condition.**

### **The first patient.**

The skin pathological process is widespread, represented by spots, blisters, milia, erosion, scales, crusts and scars, localized on the skin of the scalp, face, neck, trunk, upper and lower limbs. There are many foci of cicatricial alopecia on the skin of the scalp. The erosions are from 1 to 3 cm and are localized in the area of the forehead, nose and cervical fold, they are weep on the surface, part of the elements dry out with the formation of crusts of a mixed nature. There is a large accumulation of miliums in the area of the nasal bridge. The foci on the body are mainly in the clavicle, abdomen and lower ½ of back, in these places there are many erosions with scraps of epidermis along the edges, many of which are fresh with a bright red bottom and granulation, some at the epithelialization stage, some of the erosions are covered with serous-hemorrhagic and mixed crusts. On the skin of the lower limbs in the area of the knee joints there are multiple fresh erosions with a bright red bottom, scarring changes of pink color on the periphery. Erosion of moderate degree of exudation. Fresh blisters appear on the skin of the face, lower extremities, the tire is thin. Nail plates on the hands and feet are dystrophic or absent, slight pseudosyndactyly. Subjectively : patient feels itching, soreness.

### **The second patient.**

The skin pathological process is widespread, represented by spots, blisters, milia, erosion, scales, crusts and scars, localized on the skin of the face, trunk, upper and lower limbs. On the face in the area of the nasal dorsum, numerous grouped milia are noted against the background of an erythematous spot. The foci on the body are predominantly in the abdomen and lumbar spine, in these places there are many erosions with scraps of epidermis along the edges, many of which are fresh with a bright red bottom and granulation, some at the epithelization stage, some of the erosions are covered with serous-hemorrhagic crusts. There are white scars in the places of former erosion. The skin of the hands in the form of cicatricial atrophy, on the fingers there are single erosions with a low level of exudation, some have dried up with scraps of the epidermis along the edges. There are partial pseudosyndactyly, nail plates partially preserved on the hands. The legs are full of pseudosyndactyly, there are no nail plates. Microstomy in the oral cavity: multiple erosions on the mucosa, the tongue is almost not mobile - fusion of the frenum. Subjectively : patient feels itching, soreness.